

### Initial Patient Assessment

Assess patient's mobility levels within 8 hours of admission to the ICU and reassess at least every shift.

Start at level 1 if the patient meets any of these criteria or skip to level 2.

PaO<sub>2</sub>/FiO<sub>2</sub>

<250

Positive End-Expiratory Pressure (PEEP)

≥10 cm H₂O

O<sub>2</sub> Saturation

<90%

Respiratory Rate (RR)

Not within 10-30 per minute

Cardiac Arrhythmias or Ischemia

New Onset

Heart Rate (HR)

<60 or >120 beats per minute

Mean Arterial Pressure (MAP)

<55 or >140 mm Hg

Systolic Blood Pressure (SBP)

<90 or >180 mm Hg

Vasopressor Infusion

New or increasing

Richmond Agitation Sedation Scale (RASS)

**←**3

Riker Sedation-Agitation Scale (SAS)

<3

## Level 1: BREATHE

Patient Assessment: RASS -5 to -3; SAS 1-2

(eg, cannot participate)



### Level 2: TILT

Patient Assessment: RASS >-3; SAS >3

(eg, opens eyes; may have profound weakness)



### Level 3: SIT

Patient Assessment: RASS >-1; SAS >3

(eg, weak but may move arms/legs independently)



Patient Assessment: RASS >0; SAS >4

(eg, weak but may tolerate increased activity)



Patient Assessment: RASS >0; SAS >4

(eg, weak but may tolerate increased activity)



### **Activities**

- Maintain HOB ≥30°
- q2hr turning
- Consider continuous lateral rotation therapy (CLRT)
- Passive range of motion (ROM)2 times/day
- Up to 20° Reverse Trendelenburg/Tilt Table with lower extremity exercises/ retracting footboar Min 15 mins/Max 60 mins | 1 time/day

#### **Activities**

- Maintain HOB ≥30°
- q2hr turning
- Passive/active ROM | 3 times/day
- Up to 20° Reverse Trendelenburg/ Tilt Table with lower extremity exercises/retracting footboard Min 15 mins/Max 60 mins | 3 times/day
- Legs dependent15-20 mins | 3 times/day
- Physical therapist (PT) consultation 1 time/day

#### **Activities**

- Maintain HOB ≥30°
- q2hr turning (assisted)
- Active ROM | 3 times/day
- Encourage activities of daily living As tolerated
- Full chair position (footboard on)60 mins | 3 times/day
- Dangling, if patient can move arm against gravity | As tolerated
- PT/ Occupational Therapy (OT) actively involved | 1 time/day

#### **Activities**

- Maintain HOB ≥30°
- q2hr turning (self/assisted)
- Active ROM | 3 times/day
- Encourage activities of daily living As tolerated
- Full chair position (footboard off/feet on the floor) | 3 times/day
- Stand attempts, if patient can move leg against gravity (use a sit-to-stand lift) 3 times/day
- Pivot to chair, if tolerates partial weight bearing | 2 times/day
- PT/OT actively involved | 1 time/day

#### **Activities**

- Maintain HOB ≥30°
- q2hr turning (self/assisted)
- Active ROM | 3 times/day
- Encourage activities of daily living As tolerated
- Patient stands/bears weight >1 min 3 times/day
- Patient marches in place | 3 times/day
- Ambulate to bedside chair to achieve "out-of-bed" (use a patient lift)
  3 times/day
- PT/OT actively involved | 1 time/day

# Move to Level 2 when the Patient...

- Has acceptable oxygenation/ hemodynamics
- Tolerates q2hr turning
- Tolerates HOB >30° or up to 20° Reverse Trendelenburg

# Move to Level 3 when the Patient...

- Tolerates active-assistance exercises 2 times/day
- Tolerates lower extremity exercises against footboard/Up to 20° Reverse Trendelenburg
- Tolerates legs dependent / HOB 45°

# Move to Level 4 when the Patient...

- Tolerates increasing active exercise in bed
- Actively assists with q2hr turning or turns independently
- Tolerates full chair position 3 times/day

# Move to Level 5 when the Patient...

- Can successfully comply with all activities
- Tolerates trial periods of full chair position (footboard off/feet on the floor) 3 times/day
- Tolerates partial weight-bearing stand and pivots to chair

Continue to ambulate progressively longer distances as tolerated until patient consistently participates and moves independently.

Assessed to Level 1 or 2

**Progress to Level 2** 

**Progress to Level 3** 

**Progress to Level 4** 

**Progress to Level 5** 

**End Protocol**